



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DISCUSSION AND CORRESPONDENCE

THE POSITIVE ION IN ELECTRICAL DISCHARGE
THROUGH GASES

WHEN a metal sphere is hung upon a silk cord between the terminals of a plate glass electric machine, it will oscillate to and fro between the terminals.

When molecules of a gas are placed in a similar position, they can not behave in quite the same fashion. No one molecule can plow its way through the swarm of molecules which surround it. They are all being urged to do this. At any instant some are being urged away from the positive terminal, and some from the negative. These opposing streams of gas mingle. The collisions which result between these overcharged and undercharged molecules within such a field of force result in a continual transfer of electrical corpuscles from molecule to molecule. In such a mixture we should at any instant expect to find three classes of molecules. Those which are negatively overcharged, those which are negatively undercharged, and those which are in normal condition.

Even in open air discharge, the repelled molecules move along streamers. In particular is this the case at and near the positive terminal. Here the corpuscles and air molecules are moving in opposite directions. In rarefied gas, where the mean free path is greatly increased, these streamers become "rays."

All of the properties of these rays are in harmony with what we should expect, from our knowledge of the behavior of the metal ball and the properties of gases.

FRANCIS E. NIPHER

WASHINGTON UNIVERSITY

A NEW RECORD OF A CHESTNUT-TREE DISEASE
IN MISSISSIPPI

PROFESSOR EUGENE HILGARD, of Berkeley, Cal., told me this summer of an observation of his which is of moment to those interested in the chestnut-bark disease.

While surveying in 1856 in the northeastern part of Mississippi, he found the chestnut trees of that region, both young and old, dead.

They had been growing in a mixed forest of pine and oak and, as the other trees were in a healthy condition, were very noticeable. The dead trees were frequently of large proportions, attaining a height of 80 to 90 feet. When he saw them, these trees were beginning to decay; the bark was dropping off, leaving the trunks bare. There were no signs of insects. The region which was surveyed is a non-calcareous one.

As chestnuts are still growing in northeastern Mississippi, the epidemic which Professor Hilgard saw did not exterminate the tree in that region. It is another record of a devastating disease which the chestnut tree has endured.

Now that extra attention has been given to the chestnut and old records have been looked over, the struggle which this tree has had against attacks of fungi and of insects during the nineteenth century becomes apparent. There can hardly be a doubt but that the present range of this tree is much less extensive than formerly.

CAROLINE RUMBOLD

BLANDING'S TURTLE

TO THE EDITOR OF SCIENCE: Referring to Mr. Howe's note in SCIENCE of September 1, "Second Record for Blanding's Turtle in Concord, Mass.," and of the introduction of three pairs of the same species in Little Long Pond, Orange County, by Dr. Townsend, I beg leave to report finding this turtle at Queens, L. I., in June, 1909. It has been placed on the records of the Natural History Survey of Long Island now being made by the Brooklyn Institute of Arts and Sciences. This is the first report, so far as we know, of Blanding's turtle having been found on Long Island, but Abbott in "A Naturalist's Rambles about Home" mentions finding it in central New Jersey.

JOHN J. SCHOONHOVEN

THE MOTH OF THE COTTON WORM

TO THE EDITOR OF SCIENCE: In connection with the notices appearing in SCIENCE (October 16 and November 10) recording the occurrence far north of the moth of the cotton